High priority exotic pests of the grains industry



The following are some key high priority exotic pest threats of the grains industry. Any of these pests would have serious consequences should they enter and become established in Australia.

Karnal bunt (Tilletia indica)

OVERALL RISK - EXTREME

- Hosts are wheat, durum and triticale
- Parts of seeds are blackened and crush relatively easily
- Infected grain has a distinct fishy smell
- If it became established in Australia, access to over 45 international markets would be restricted and grain price would be significantly reduced



aDIL

Khapra beetle (Trogoderma granarium)

OVERALL RISK - HIGH

- Adults are small (2-3 mm long) and do not fly
- Spread in infested grain
- Larvae are hairy and can survive for over a year without food
- Phosphine fumigation gives poor control
- If established, it would affect market access



viinistry of Agriculture and Regional Development

Phosphine resistant strains of stored grain insects

OVERALL RISK - MEDIUM

- Stored grain insects with strong resistance to phosphine have been detected internationally and in Australia
- Exports are threatened due to insects surviving in stored grain
- Incorrect phosphine fumigation and poor grain storage practices increase the selection of resistant insects
- Live insects remaining after fumigation should be reported and tested for resistance



Hessian fly and barley stem gall midge (Mayetiola destructor and M. hordei)

OVERALL RISK - HIGH

- Adults are small (2-4 mm long) and look like mosquitoes
- Pupae have a 'flaxseed' appearance
- Attack leaves, stems and heads of cereals
- Most chemical controls are not effective
- Cereal crop losses up to 40% could occur



scott Bauer, USDA Agricultural Res Service, Bugwood

High priority exotic pests of the grains industry



Sunn pest (Eurygaster integriceps)

OVERALL RISK - MEDIUM

- Brown bug with wide oval-shaped body (12 mm long) with a wide triangular head
- Attacks most cereal crops
- Colonies can be seen on cereal heads in spring
- Injects enzymes into the plant as it feeds which can result in grain damage and abortion



Barley stripe rust (Puccinia striiformis f. sp. hordei)

OVERALL RISK - HIGH

- Would infect barley in all Australian growing regions
- Approximately 80% of Australia's barley varieties would be susceptible
- Yellow stripes of fungal spores produced between veins of leaves
- Can be spread by wind and rain, or on clothing, machinery and tools
- Any stripe rust on barley should be reported



Pat Hayes, Oreg

Wheat stem rust, pathotype Ug99 (Puccinia graminis f. sp. tritici)

OVERALL RISK - HIGH

- Pathotype identified in Uganda in 1999 that has overcome several stem rust resistance genes
- Many Australian wheat varieties will be susceptible
- Elliptical blisters produced on stems, which break open to reveal a mass of rust coloured spores
- Stem rust on known resistant varieties should be reported



inistry of Agriculture and egional Development

About the Grains Farm Biosecurity Program

The Grains Farm Biosecurity Program is managed by PHA and funded by growers through Grain Producers Australia together with the governments of New South Wales, Queensland, South Australia, Victoria and Western Australia. Grains Biosecurity Officers in these five states develop and deliver materials to raise awareness and deliver training to growers, consultants and other industry stakeholders. Go to **grainsbiosecurity.com.au** for more information.

If you see anything unusual, call the Exotic Plant Pest Hotline



Disclaimer: The material in this publication is for general information only and no person should act, or fail to act on the basis of this material without first obtaining professional advice. Plant Health Australia and all persons acting for Plant Health Australia expressly disclaim liability with respect to anything done in reliance on this publication.

