# Barley stripe rust



## What is barley stripe rust?

Barley stripe rust (*Puccinia striiformis* f. sp. *hordel*) is a fungal pest of barley, potentially affecting crops in all Australian growing regions. Barley stripe rust would have serious economic impacts following establishment in Australia, as it is estimated that approximately 60% of the barley varieties grown here would be susceptible. Damage to barley plants varies depending on the plant growth stage. Crop losses due to barley stripe rust are typically around 10%, but in susceptible varieties they may be up to 100%.

#### What does it look like?

The fungus produces stripes of rust pustules between the veins of leaves, and can also form on barley heads. The pustules may be more yellow than orange, and hence the disease is sometimes referred to as yellow rust. Barley stripe rust can build up rapidly if conditions are cool and wet, and infection is often first noticed as 'hot spots' within the crop.

#### What can it be confused with?

Barley grass stripe rust (present in the eastern states of Australia), and occasionally wheat stripe rust, can infect barley. However, any barley plants showing stripe rust symptoms should be sent for identification through laboratory analysis.

### What should I look for?

The development of stripe rust symptoms on any barley plants. In certain conditions 'hot spots' of symptoms can develop in the crop.

#### How does it spread?

Rust spores are small, light and may survive for several days. They can be spread over large distances by wind and easily attach to clothing, machinery and tools, allowing movement and spread between farms and regions. They may also transfer across the world on travellers' clothing.




Stripe rust forms yellow orange stripes along leaf veins

Lee F. Jackson, UC Statewide IPM Program



Stripe rust symptoms can develop as 'hot spots' in the crop

John Francis



The stripes of stripe rust are made up of many tiny pustules arranged between the leaf veins

Jack Kelly Clark, UC Statewide IPM Program



#### Where is it now?

Barley stripe rust occurs in Asia, Europe, Central Africa and North, Central and South America.

## How can I protect my farm from barley stripe rust?

Check your farm frequently for the presence of new pests and unusual symptoms. Make sure you are familiar with common grain pests so you can tell if you see something different.

If you see anything unusual, call the **Exotic Plant Pest** Hotline on 1800 084 881.

#### GRAINS FARM BIOSECURITY PROGRAM

An initiative of Plant Health Australia and Grain Producers Australia





IF YOU SEE ANYTHING UNUSUAL, CALL THE EXOTIC PLANT PEST HOTLINE

1800 084 881



Stripe rust can also show symptoms on spikes, as demonstrated by this Wheat stripe rust infection

Colin Wellings



Leaves take on a yellow colour under high levels of infection

PaDIL



Advanced symptoms present on barley leaf sheaths

Darrell Wesenberg, Bugwood.org

DISCLAIMER: The material in this publication is for general information only and no person should act, or fail to act on the basis of this material without first obtaining professional advice. Plant Health Australia and all persons acting for Plant Health Australia expressly disclaim liability with respect to anything done in reliance on this publication.